

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

81794

(19)



JAPANESE PATENT OFFICE

PATENT ABSTRACTS OF JAPAN

(11) Publication number: 06213888 A

(43) Date of publication of application: 05 . 08 . 94

(51) Int. Cl

G01N 33/53
G01N 33/549

(21) Application number: 04361873

(22) Date of filing: 24 . 12 . 92

(71) Applicant: FUJI YAKUHI KOGYO KK

(72) Inventor:
OKADA YASUNORI
FUJIMOTO NOBORU
MORI NOBUKO
OUCHI EIKO
SAKAI CHIE
TOKAI HIDEAKI
HAYAKAWA TARO
IWATA KAZUSHI

(54) IMMUNOLOGICAL DETERMINATION METHOD OF
HUMAN 72-KDA GELATINASE/IV-TYPE
COLLAGENASE

(57) Abstract:

PURPOSE: To determine human pro MMP-2 within a sample to be tested rapidly with high sensitivity and accuracy by using two types of monoclonal antibodies which specifically combines with human 72-kDa gelatinase/IV-type collagenase (human pro MMP-2).

CONSTITUTION: Two types of monoclonal antibodies which specifically combine with human pro-MMP-2 are used as carriers which combine with solid phase carriers

or antibodies which give labeled substances, thus performing measurement immunologically by the sandwich method. Balls, microplates, sticks, fine particles, or test tubes made of for example polystyrene and polycarbonate for well adsorbing protein such as antibodies can be selected arbitrarily and used as the solid phase carriers. On the other hand, after an object containing antibodies is fractionated with ammonium sulfate as antibodies giving the labeled substance, anion exchange gel and IgG fraction are performed and further specifically combined part Fab' which is obtained by reduction after pepsin digestion can be used.

COPYRIGHT: (C)1994,JPO&Japio